

POWERS OF M_{24} -TWISTED SIEGEL PRODUCT EXPANSIONS ARE MODULAR

MARTIN RAUM

ABSTRACT. We prove that several product expansions suggested by Cheng and Duncan are Siegel modular forms. They arising from twists of elliptic genera of symmetric powers of $K3$ surfaces. Our result shows that they can be represented as products of rescaled Borcherds products.

1. INTRODUCTION

In [EOT10], Eguchi, Ooguri, and Tachikawa suggested that Fourier coefficients of the elliptic genus of $K3$ surfaces have a natural decomposition in terms of the dimensions of the Mathieu group M_{24} —see [GV12] for a discussion of this decomposition in the light of more recent results. This observation has led to a intense study of a possible M_{24} moonshine by both physicists and mathematicians. Cheng suggested the construction of M_{24} twisted elliptic genera for symmetric powers of $K3$ surfaces [Che10]. These lead to product expansions Φ_g , where g is a conjugacy class in M_{24} . In a recent paper, Cheng and Duncan [CD12] discussed this idea in more detail and made the following conjecture (see Section 3 for details on the notation):

Conjecture 1.1 ([CD12]). *For all conjugacy classes g of M_{24} , the product $\Phi_g(Z)$ is a Siegel modular form with character for the Siegel type congruence subgroup $\Gamma_0^{(2)}(N_g)$. The conjectured levels N_g is given in Table 1 or [CD12, Table 1].*

To the author's knowledge, so far, only the cases $g \in \{2A, 3A, 4B\}$ were solved completely. We are able to resolve several new cases of this conjecture. More precisely, we prove the following result:

Theorem 1.2. *For all g in Table 1, $\Phi_g^{p_g}$ is a Siegel modular form with character for $\Gamma_0^{(2)}(\tilde{N}_g)$ and weight $p_g k_g$.*

g	2A	2B	3A	3B	4A	4B	4C	5A	7AB	8A	11A	23AB
n_g	2	2	3	3	4	4	4	5	7	8	11	23
N_g	2	4	3	9	8	4	16	5	7	8	11	23
$\tilde{N}_g$	2	4	3	9	8	4	16	5	7	16	121	529
k_g	6	4	4	2	2	3	1	2	1	1/2	0	-61/18
p_g	3	1	2	3	6	6	4	3	4	12	6	132

TABLE 1. Conjectured and proved levels of Φ_g .

A more detailed statement is given in Theorem 4.1, of which the above theorem is a weak version. The proof of the later is largely computer assisted.

Note that the p_g are not necessarily optimally, a fact which stems from an ambiguity in the arising system of linear equations. For example, it is known that p_{2A} can be chosen to be 1, but in Table 1 we only give $p_{2A} = 3$. It should also be mentioned that the weight of Φ_g for $g = 23AB$ is curious.

While so far in most papers on product expansions in string theory only Borcherds products have been taken into account, we include rescaled Borcherds products into our consideration—see Section 2, and in particular the last remark in said section, for a discussion of this construction. Seemingly the new cases that we treat are not Borcherds products, and thus do not fall under the scope of considerations in the literature.

2000 *Mathematics Subject Classification.* Primary 11F46, 81T30; Secondary 11-04.

Key words and phrases. Mathieu Moonshine, Siegel product expansions, twisted elliptic genera of $K3$ surfaces.

The author holds a scholarship from the Max Planck society.

We proved that in all remaining cases the Φ_g are not rescaled Borcherds products of level N_g , but the author is optimistic that they are for large enough $\tilde{N}_g$ in the sense of Theorem 1.2. The lack of sufficient computational power, prevented the author from checking such $\tilde{N}_g$.

The paper is organized as follows. In Section 2, we recall the theory of Borcherds products that we will make use of. We also introduce rescaled Borcherds products. Section 3 contains a revision of the material in [CD12] that is relevant to this paper. The proof of Theorem 4.1 is discussed in Section 4.

2. BORCHERDS PRODUCTS AND RESCALED BORCHERDS PRODUCTS

Given $\Gamma \subseteq \mathrm{SL}_2(\mathbb{Z})$, we write $J_{k,m}^!(\Gamma)$ for the space of weakly holomorphic Jacobi forms of weight k and index m for $\Gamma \ltimes \mathbb{Z}^2$ —see [EZ85] for details. The set of cusp classes of Γ will be denoted by $\mathcal{C}(\Gamma)$, and for any cusp class $\mathfrak{c} \in \mathcal{C}(\Gamma)$, we write $h_{\mathfrak{c}}(\Gamma)$ and $e_{\mathfrak{c}}(\Gamma)$ for the width and the denominator of $\mathfrak{c}$, respectively. If $\Gamma = \Gamma_0(N)$ for some positive integer N , then we write $h_{\mathfrak{c}}(N)$ and $e_{\mathfrak{c}}(N)$ instead. We set $N_{\mathfrak{c}}(N) = Ne_{\mathfrak{c}}^{-1}$, and also suppress the argument N , whenever it is clear for the context.

According to [CG08], if $\Gamma = \Gamma_0(N)$, we choose the representatives fe^{-1} for the cusps, where $e \geq 1$, $e \mid N$. Given $\phi \in J_{k,m}^!(\Gamma_0(N))$ and $\mathfrak{c} \in \mathcal{C}(\Gamma)$, we write $\phi_{\mathfrak{c}}$ for the Fourier expansion of ϕ at that representative of $\mathfrak{c}$.

Siegel modular forms (of genus 2) are certain functions on the Siegel upper half space

$$\mathbb{H}_2 := \{Z = X + iY \in \mathrm{M}_2(\mathbb{C}) : Z = Z^T, Y \text{ is positive definite}\}.$$

In order to give a precise definition, write $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ for a typical element $\gamma \in \mathrm{Sp}_2(\mathbb{R})$ and let the weight k Siegel slash action be

$$(f|_k \gamma)(Z) = \det(CZ + D)^{-k} f((AZ + B)(CZ + D)^{-1}).$$

Definition 2.1. Let $f : \mathbb{H}_2 \rightarrow \mathbb{C}$ be a holomorphic function. We call f a Siegel modular form of weight k with character χ for $\Gamma \subset \mathrm{Sp}_2(\mathbb{Z})$ if and only if $\chi(\gamma) f|_k \gamma = f$ for all $\gamma \in \Gamma$.

We write $[\Gamma, k, \chi]$ for the space of such Siegel modular forms.

We refer the reader to [Fre83] for details on Siegel modular forms.

Write $Z = \begin{pmatrix} \tau_1 & z \\ z & \tau_2 \end{pmatrix}$, and set $q_1 = \exp(2\pi i \tau_1)$, $\zeta = \exp(2\pi i z)$, and $q_2 = \exp(2\pi i \tau_2)$. Siegel modular forms for suitable subgroups of $\mathrm{Sp}_2(\mathbb{Z})$, including congruence subgroups $\Gamma_0^{(2)}(N)$ of Siegel type, have a Fourier expansion

$$\sum_{(n,r,m)} a(n,r,m) q_1^n \zeta^r q_2^m.$$

The sum runs over triples (n, r, m) of integers.

In this paper, we put our focus on product expansions of Siegel modular forms. We start by recalling the following special case of a theorem by Clery and Gritsenko [CG08], which is a refinement and reformulation in terms of Jacobi forms, of Borcherds's result [Bor98] on product expansions for Siegel modular forms. In order to state it, note that the Fourier expansion of index 1 Jacobi forms, can be by discriminants $4n - r^2$ only.

Theorem 2.2 (Clery-Gritsenko). Let $\phi(\tau, z) = \sum_{n,r} c(4n - r^2) q^n \zeta^r \in J_{0,1}^!(\Gamma_0(N))$. Write $c_{\mathfrak{c}}$ for the Fourier coefficients of $\phi_{\mathfrak{c}}$, and assume that for all cusp classes $\mathfrak{c} \in \mathcal{C}(\Gamma_0(N))$ we have that $h_{\mathfrak{c}} N_{\mathfrak{c}}^{-1} c_{\mathfrak{c}}(D) \in \mathbb{Z}$, if $D \leq 0$. Let

$$B[\phi](Z) := q_1^{e_{q_1}} \zeta^{e_{\zeta}} q_2^{e_{q_2}} \prod_{\mathfrak{c} \in \mathcal{C}(\Gamma_0(N))} \prod_{(n,r,m) > 0} (1 - (q_1^n \zeta^r q_2^m)^{N_{\mathfrak{c}}})^{h_{\mathfrak{c}} N_{\mathfrak{c}}^{-1} c_{\mathfrak{c}}(4nm - r^2)}$$

where

$$e_{q_1}[\phi] := \frac{1}{24} \sum_{\mathfrak{c} \in \mathcal{C}(\Gamma_0(N)), l \in \mathbb{Z}} h_{\mathfrak{c}} c_{\mathfrak{c}}(-l^2), \quad e_{\zeta}[\phi] := \frac{1}{2} \sum_{\mathfrak{c} \in \mathcal{C}(\Gamma_0(N)), l > 0} l h_{\mathfrak{c}} c_{\mathfrak{c}}(-l^2), \quad e_{q_2}[\phi] := \frac{1}{4} \sum_{\mathfrak{c} \in \mathcal{C}(\Gamma_0(N)), l \in \mathbb{Z}} l^2 h_{\mathfrak{c}} c_{\mathfrak{c}}(-l^2).$$

Set

$$k[\phi] := \frac{1}{2} \sum_{\mathfrak{c} \in \mathcal{C}(\Gamma_0(N))} h_{\mathfrak{c}} N_{\mathfrak{c}}^{-1} c_{\mathfrak{c}}(0).$$

Then $B[\phi]$ is a meromorphic Siegel modular form of weight $k[\phi]$ with character for $\Gamma_0^{(2)}(N)$. If $c_{\mathfrak{c}}(D) \geq 0$ for all $D < 0$, then $B[\phi]$ is holomorphic.

We will analyze product expansions of the form

$$(2.1) \quad \prod_i B[\phi_i](n_i Z),$$

where the product is finite and the ϕ_i are weakly holomorphic Jacobi forms of weight 0 and index 1 for $\Gamma_0(N_i) \ltimes \mathbb{Z}^2$. The n_i are positive integers that can be chosen freely. By the above theorem, (2.1) is a meromorphic Siegel modular form with character for $\Gamma_0^{(2)}(\text{lcm}\{N_i n_i\}_i)$.

The rescaled Borcherds product (2.1) has the following product expansion:

$$(2.2) \quad \prod_i B[\phi_i](n_i Z) = q_1^{\sum_\phi n_i e_{q_1}[\phi_i]} \zeta^{\sum_\phi n_\phi e_\zeta[\phi_i]} q_2^{\sum_\phi n_\phi e_{q_2}[\phi_i]} \prod_{d \mid N} \prod_{(n,r,m) > 0} (1 - (q_1^n \zeta^r q_2^m)^d)^{c_d(4nm-r^2)},$$

where $N = \text{lcm}\{N_\epsilon(N_i) n_i\}_{i, \epsilon \in \mathcal{C}(\Gamma_0(N_i))}$ and the $c_d(D)$ are the Fourier coefficients of the following Jacobi forms:

$$(2.3) \quad \psi_d(\tau, z) = \sum_i \sum_{\epsilon \in \mathcal{C}(\Gamma_0(N_i)), N_\epsilon(N_i) = d/n_i} \frac{h_\epsilon(N_i)}{N_\epsilon(N_i)} \pi_{\text{FE}}((\phi_i)_\epsilon)(\tau, z).$$

Here, π_{FE} is the projection of the Fourier expansion to those terms with integral exponents:

$$\pi_{\text{FE}}\left(\sum_{n \in \mathbb{Q}} a(n) q^n\right) := \sum_{n \in \mathbb{Z}} a(n) q^n.$$

It is not hard to see that ϕ_d is a Jacobi form of weight 0 and index 1 for the Jacobi group $\Gamma_0(\text{lcm}\{N_i\}_i) \ltimes \mathbb{Z}^2$.

3. SIEGEL PRODUCT EXPANSIONS BY CHENG-DUNCAN

Consider the following family of weak Jacobi forms $\mathcal{Z}_g$, the M_{24} -twisted elliptic genus.

$$\mathcal{Z}_g(\tau, z) := \chi(g) \frac{\phi_{0,1}(\tau, z)}{12} + \tilde{T}_g(\tau) \phi_{-2,1}(\tau, z),$$

where $\tilde{T}_g$ is given in [CD12, Table 2]. By [CD12, Proposition 3.1], the $\mathcal{Z}_g$ are weak Jacobi forms of weight 0 and index 1 for the Jacobi group $\Gamma_0(N_g) \ltimes \mathbb{Z}^2$. We write

$$(3.1) \quad \mathcal{Z}_g(\tau, z) = \sum_{n \geq 0, r \in \mathbb{Z}} c_g(4n - r^2) q^n \zeta^r$$

for the Fourier expansion of $\mathcal{Z}_g$.

In [CD12, Section 4], the following product expansion was given

$$\Phi_g(Z) := q_1 \zeta q_2 \prod_{(n,r,m) > 0} \exp\left(-\sum_{k=1}^{\infty} \frac{c_g(4nm-r^2)}{k} (q_1^n \zeta^r q_2^m)^k\right).$$

The product runs over triples of integers (n, r, m) , and $(n, r, m) > 0$ means that $n > 0$, or $n = 0$ and $m > 0$, or $n = m = 0$ and $r < 0$.

In analogy with [Che10, (3.9)], we can rewrite this as

$$(3.2) \quad \Phi_g(Z) = q_1 \zeta q_2 \prod_{d \mid n_g} \prod_{(n,r,m) > 0} (1 - (q_1^n \zeta^r q_2^m)^d)^{c_{g,d}(4nm-r^2)},$$

where

$$c_{g,d}(D) := -d^{-1} \sum_{d' \mid d} \mu(d/d') c_{g,d'}(D).$$

The Möbius function is denoted by μ .

The $c_{g,d}$ are Fourier coefficients of a weak Jacobi form of weight 0 and index 1, as can be seen as follows:

$$(3.3) \quad \mathcal{Z}_{g,d}(\tau, z) = \sum_{n \geq 0, r \in \mathbb{Z}} c_{g,d}(4n - r^2) q^n \zeta^r := -d^{-1} \sum_{d' \mid d} \mu(d/d') \mathcal{Z}_{g,d'}(\tau, z).$$

4. RESULTS

Theorem 4.1. *For all g in Table 1, $\Phi_g^{p_g}$ is of the form $\prod_i B[\phi_i](n_i Z)$. The ϕ_i and n_i are given as Sage readable data in “sol_ $\mathbf{g}$.sobj”, where $\mathbf{g}$ must be replaced by the corresponding label of the conjugacy class.*

Proof of Theorem 4.1. The proof is largely computer assisted. Fix g and recall the product expansion (3.2) for Φ_g . In order to prove the theorem, it is sufficient to find Jacobi forms ϕ_i of weight 0 and index 1 such that $\mathcal{Z}_{g,d}$ defined in (3.3) equals ψ_d defined in (2.3)

In order to find such forms one can fix a basis of the space of weight 0 and index 1 weak Jacobi forms for $\Gamma_0(\tilde{N}_g) \ltimes \mathbb{Z}^2$. Computing the Borcherds lift of each basis element leads to a product expansion like in (2.1). Comparing the Jacobi forms ψ_d in Equation (2.3) and $\mathcal{Z}_{g,d}$ in Equation (3.3) leads to a system of linear equations that can then be solved.

For each g , we give ϕ_i ’s that work in the form $f_0 \phi_{0,1}(\tau, z)/12 + f_2(\tau) \phi_{-2,1}(\tau, z)$, where f_0 is a constant and $f_2(\tau)$ is a sum of eta products. Note that the Fourier expansion of ϕ_c can be computed from $(f_2)_c$. The data is stored in the files “sol_ $\mathbf{g}$.sobj”, where $\mathbf{g}$ must be replaced by the label of a conjugacy class in Table 1. The scripts used to compute this data is published at [Rau]. $\square$

Acknowledgement: The author thanks Kathrin Bringmann for helpful conversations about the product expansions in [CD12].

REFERENCES

- [Bor98] R. Borcherds, *Automorphic forms with singularities on Grassmannians*, Invent. Math. **132** (1998), no. 3, 491–562.
- [CD12] M. Cheng and J. Duncan, *The Largest Mathieu Group and (Mock) Automorphic Forms*, ArXiv: 1201.4140v1, 2012.
- [CG08] F. Cléry and V. Gritsenko, *The Siegel modular forms of genus 2 with the simplest divisor*, ArXiv: 0812.3962v1, 2008.
- [Che10] M. Cheng, *K3 Surfaces, $\mathcal{N} = 4$ Dyons, and the Mathieu Group M_{24}* , ArXiv: 1005.5415v2, 2010.
- [EOT10] T. Eguchi, H. Ooguri, and Y. Tachikawa, *Notes on the K3 Surface and the Mathieu group M_{24}* , ArXiv: 1004.0956, 2010.
- [EZ85] M. Eichler and D. Zagier, *The Theory of Jacobi Forms*, Birkhäuser, Boston, 1985.
- [Fre83] E. Freitag, *Siegelsche Modulformen*, Grundlehren der mathematischen Wissenschaften, Springer Verlag, Berlin, 1983.
- [GV12] M. Gaberdiel and R. Volpato, *Mathieu Moonshine and Orbifold K3s*, ArXiv: 1206.5143, 2012.
- [Rau] M. Raum, *Homepage*, <http://people.mpim-bonn.mpg.de/mraum/xhtmll/index.xhtmll>.

MAX PLANCK INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY
E-mail address: MRaum@mpim-bonn.mpg.de